




Interaction of Asian Monsoon and global oceans

—in a changing climate



Interaction of Asian Monsoon and global oceans

1. Past: A bit of Monsoon history

2. Present: The Monsoon we know/live

- Role of Ocean on the Monsoon
 - Interannual variability of the Monsoon – ENSO, IOD, etc.
 - Intraseasonal variability of the Monsoon
 - SST–convection relationship




3. Trends/Future: Changing Oceans and the Monsoon

- Changing ENSO–Monsoon interactions
- Indian Ocean warming
- Atlantic warming
- Interplay between the tropical oceans

4. Monsoon Modeling

History of Monsoon, 80 Mya


Plate tectonics – factors leading to modern monsoon



Indian plate motion: rate of $\sim 15\text{cm/yr}$
– collided with Eurasia c.20–30 Mya.
Tethys evolved into the present day Indian Ocean


History of Monsoon, 80 Mya

Remnants of Deccan Volcanism




Rise of the Himalayas and strengthening of the monsoon

Geography



30Mya

Rainfall





The Himalaya–Tibetan plateau uplift during late Tertiary strengthened the Asian Monsoon.

Confinement of African monsoon to a thin band, expanding the subtropical desert is also seen.



Role of Orography:

1. Upward deflection of large scale horizontal flow by orography
2. Uplifted moist air will expand, cool and condense – forming clouds


10Mya



Present




Monsoon weakening in NW and collapse of Indus Valley




Variation in the timing and intensity of the monsoon affects lake-water Oxygen isotope and alters relative hydrologic balance between evaporation and precipitation in the lake.

Presence/absence of ostracods confirms this.



History of Monsoon, 1500 AD

Trade Winds – History Changers!






Using the trade wind secret, Indian/Asian and Arabian traders were able to dominate the lucrative market by concealing the true source of their cargoes for centuries.

Mediterranean sailors from Egypt and Europe were fearful of open-sea sailing and their ships hugged the coastlines

– until 1st century BC when a stranded sailor revealed the secret to Egyptian Officials.


Monsoon, manifested by Winds

Trade Winds




Monsoon, manifested by Surface Temperatures

Seasonal migration of solar insolation




← January

July →




Monsoon, manifested by Thermal Contrast

Land and sea warms at different rates, sp.heat capacity



Monsoon, manifested by Pressure Gradient


Monsoon Trough and 120° shift in wind direction



Madagascar–Mascarene High/Anticyclone and Monsoon Trough



Monsoon, manifested by conducive SST

SST > 28°C conducive for enhanced convective activity



Monsoon, manifested by Rainfall

Seasonal Migration of Tropical Convergence Zone (TCZ)




Annual evolution of daily mean Winds at 850 hPa and Precipitation

Interannual Variability of the Monsoon


predictable mode analysis

ocean-atmospheric processes – 47% of total variance

EOFs and corresponding PCs of precipitation






Correlation of SST with PC



Interannual Variability of the Monsoon – ENSO


El Niño Southern Oscillation (ENSO) is a major driver

Generally, weak monsoon coincides with El Niño and strong monsoon with La Niña.





Interannual Variability of the Monsoon - ENSO

ENSO – neutral conditions – Bjerknes feedback



Interannual Variability of the Monsoon – ENSO


ENSO – neutral conditions – Bjerknes feedback



- Strong east–west asymmetry over the equatorial Pacific: Walker circulation (with easterly winds at the surface) in the atmosphere and a 5degC SST contrast between west warm pool and east cold tongue.
- Under easterly winds, surface ocean currents flow poleward a few degrees away from the equator following the Ekman dynamics → upwells water into the surface layer on the equator.
- If the thermocline is close to surface, this equatorial upwelling brings cold thermocline water into the mixed layer, causing an SST cooling. On the equator where the Coriolis force vanishes, surface currents flow in the wind direction, resulting in the westward South Equatorial Current (SEC). The equatorial upwelling and the SEC shoals the thermocline in the east and deepens it in the west.
- On the other hand, the eastward SST cooling limits deep convection to the west and maintains a pressure gradient that drives the easterly winds along the equator. This circular argument indicates that ocean–atmosphere interaction is at a state of positive feedback.

Interannual Variability of the Monsoon – ENSO

Bjerknes feedback – positive feedbacks couple thermocline slope, SST, zonal winds



The background appears stable, but because it is maintained by positive feedbacks, weakening any element weakens the entire system (Bjerknes 1966).

An example is westerly winds in the west originating in the MJO. Though these may be transient, they can result in longer-standing disruption of the background, and a coupled chain reaction leading to El Niño.


While trades and upwelling may continue in the east, events in the west can deepen the thermocline in the east, warming SST remotely there (Wyrtki 1975).

More about westerly wind bursts: Chen et al. *Nature Geoscience*, 2015

Interannual Variability of the Monsoon – ENSO


Walker Circulation

Neutral / La Nina conditions




Gilbert Walker

El Nino conditions





Interannual Variability of the Monsoon – ENSO

Kelvin Waves



Downwelling phase: The thick warm layer sloshes east, pushing down the thermocline as it goes, thus we call this a “downwelling” wave. The thermocline is the boundary between the warmer, near surface mixed layer and colder deeper water. Because of this downward push as the wave travels eastward, it is harder for the colder, deeper water to affect the surface so near-surface temperatures are often above average. This will often (not always) warm the surface temperatures and plant the seeds for an El Niño.

El Niños are “usually” followed by La Niñas




Interannual Variability of the Monsoon – ENSO

El Niños are “usually” followed by La Niñas

Low-level westerly wind anomalies, not only trigger eastward-moving oceanic Kelvin waves at the equator, but also westward-moving waves just north and south of the equator (called Rossby waves). While Kelvin waves are pushing warm water east, these Rossby waves move cooler subsurface water toward the west. They then bounce off the western side of the tropical Pacific (around Indonesia) and have a return trip, traveling eastward near the equator.


On their eastward trip, these waves also promote cooler water, and can neutralize or reverse El Niño around 6 months after the westerly wind bursts. This cool pulse interrupts the positive feedback mechanism responsible for the growth of an El Niño, ending El Niño and promoting La Niña development.

Since stronger El Niño events often involve stronger westerly wind anomalies, these events tend to trigger stronger Rossby waves and stronger tendencies for El Niño to decay and possibly reverse after peaking at the end of a calendar year.





Interannual Variability of the Monsoon – ENSO

but not all El Niño years are drought years – 50%



Other flavors of El Nino?
Other tropical oceans?
Or... does monsoon has its
own identity?




Interannual Variability of the Monsoon


predictable mode analysis

ocean-atmospheric processes – 47% of total variance

EOFs and corresponding PCs of precipitation



Correlation of SST with PC




Interannual Variability of the Monsoon – IOD

Indian Ocean Dipole (IOD)

aka Indian Ocean Zonal Mode (IOZM)

aka the controversial mode



DIPOLES, TEMPERATURE GRADIENTS, AND TROPICAL CLIMATE ANOMALIES

BY STEFAN HASTENRATH

An examination into misleading terminology.

Comment on “Dipoles, Temperature Gradients, and Tropical Climate Anomalies”

—RAGHU MURTUGUDDE AND
ANTONIO J. BUSALACCHI

Earth System Science Interdisciplinary Center,
University of Maryland, College Park,
College Park, Maryland


—JULIAN P. MCCREARY JR.

International Pacific Research Center,
University of Hawaii, at Manoa, Honolulu, Hawaii

Interannual Variability of the Monsoon – IOD


IOD enhances local meridional circulation – monsoon rains

Temporal evolution of IOD →



Interannual Variability of the Monsoon


Internal variability?



Intraseasonal variability of the Monsoon

Monsoon is characterized by large amplitude intraseasonal oscillations (ISO) –active and break spells of rain.


They represent a very large signal and hence potentially predictable




Intraseasonal variability of the Monsoon

— MISO, BSISO... different names... MJO??

MISO: Monsoon Intraseasonal Oscillation
e.g. Suhas et al. 2013 – Rainfall anomalies



BSISO: Boreal Summer Intraseasonal Oscillation
e.g. Lee et al. 2013 – OLR/Wind (850 hPa) anom.




1. Northward propagation
2. Ocean-atmosphere coupled interaction
3. Arabian Sea – Bay of Bengal – South China Sea

Intraseasonal variability of the Monsoon


— contribution to the seasonal mean

Higher frequency of occurrence of active conditions would result in stronger than normal seasonal mean, and vice-versa.




Spatial structure of the summer ISOs have certain similarity with that of the summer seasonal mean.

Active – Break composite




Strong – Weak monsoon composite




Intraseasonal variability of the Monsoon

— SST variability over the monsoon basins




SST & heat flux anomalies associated with monsoon ISV are observed over a large domain, Arabian Sea → Bay of Bengal → South China Sea → w. North Pacific



Intraseasonal variability of the Monsoon

— Source of SST variability

Intraseasonal SST over monsoon basins: driven by **Latent heat flux** (LHF is dominant due to stronger winds) and **Shortwave flux** (SWF) anomalies.



SST tendency equation:

$$\frac{\partial T_s}{\partial t} = \frac{F_{\text{tot}}}{\rho c_p \text{MLD}}$$


Quantitatively:

F_{tot} of 50 Wm^{-2} , MLD of 40 m, standard ρ & $c_p \Rightarrow F_{\text{tot}} / \rho c_p \text{MLD} = 0.025^\circ\text{C day}^{-1}$

SST change of 0.8°C in 40 days $\Rightarrow \partial T_s / \partial t = 0.02^\circ\text{C day}^{-1}$

Intraseasonal variability of the Monsoon


— Role of SST variability on the Monsoon ISV



positive SST anomalies

⇒ destabilize lower atmospheric column

⇒ convective activity



Intraseasonal variability of the Monsoon

— Mixed layer is important

$$\frac{\partial T_s}{\partial t} = \frac{F_{\text{tot}}}{\rho c_p * \text{MLD}}$$

Comparison of observed and model (CFSv2) ISV


ISV is overestimated in the model.
Where is it coming from?

Flux Contribution => SST Tendency

The increased SST anomalies in the model are **comparable** to the simulated net surface flux anomalies, For 30 W m^{-2} (30m MLD), $dT = 0.025\text{C day}^{-1}$.

Wind Contribution => LHF

Using the bulk aerodynamic equations, an overestimation of 1ms^{-1} of wind speed is **comparable** to an increase of 14 W m^{-2} of latent heat flux anomalies, in the model.




Flux Contribution => SST Tendency

Wind Contribution => LHF

Intraseasonal variability of the Monsoon

— Mixed layer is important

JJAS MLD Diff. [CFSv2 - Boyer]




Biases in ISV of SST correlated with biases in ISV of MLD

For the same magnitude of fluxes, change in SST is different:


Shallow MLD \rightarrow ISV amplified

Deep MLD \rightarrow ISV weakened

$r = 0.5$, significant at 95% levels




Intraseasonal variability of the Monsoon



SST- precipitation relationship

— Lower and Upper Threshold for convection?



SSTs above **26 °C** conducive for increased convection

Are SSTs important for convection, beyond **28.5 °C** ?

SST-precipitation relationship

— Earlier studies: Upper Threshold and CAPE




Figure 1. SST-convective relation in the global tropics (reproduced from Waliser *et al.*, 1993) for monthly values in $2^\circ \times 2^\circ$ latitude-longitude squares of global tropics 25°S to 25°N for the period 1975–1985: (a) SST and HRC and (b) SST and OLR. The right ver-




FIG. 3. OLR-SST (W m^{-2} and $^\circ\text{C}$ with ordinate scale reversed) pairs for (a) the tropical Pacific Ocean in the region 5°S – 5°N , 120°E – 90°W , (b) the northern Indian Ocean for the region equator– 20°N , 55° – 80°E , and (c) the northern Indian Ocean, equator– 20°N , 80° – 100°E . The geographical areas are depicted by the areas within the shaded boxes on map panel. Mean monthly values are plotted for Mar, Apr, and May for all points in the boxes. Data are from 1972–89.


Earlier studies: Precipitation increases monotonously at SSTs beyond 26°C , but limited to: **Upper threshold of $28 - 29^\circ\text{C}$**

Explanation given: Precipitation tends to occur where positive convective available potential energy (CAPE) exists

→ the occurrence of deep convection will tend to squelch CAPE?

SST- precipitation relationship

- negative relationship over west Pacific, at SSTs > 29°C?




SST- precipitation relationship

Clausius-Clapeyron

— moisture increases with increasing temperature

The Clausius–Clapeyron relation implies that specific humidity and hence atmospheric moisture would increase roughly exponentially with temperature –

7%/°C – substantially smaller than the sensitivity change documented.




C-C also controls how
“wet gets wetter, and dry gets drier”.

Observations suggest that precipitation and total atmospheric water have increased at about the same rate over past 2 decades.


→ with SST, moisture has increased

→ with moisture, rainfall has increased




SST- precipitation relationship

— Does high SSTs have an active role on Monsoon?



SSTs averaged over the open basins





mean SSTs are above 28.5°C

SST-precipitation relationship

— it's a lead-lag relationship!


The SST-precipitation relationship have different **lead-lags** over the Arabian Sea and the Bay of Bengal/South China Sea



The magnitude of the correlation refers to the intensity of the driving force, and the corresponding lag (lead) time denotes how quickly the atmosphere responds to the SST anomalies and vice versa.



SST- precipitation relationship

lags considered, 1°C rise in SST \rightarrow 2 mm/day in rainfall





SST- precipitation relationship

lags considered, 1°C rise in SST \rightarrow 2 mm/day in rainfall




SST- precipitation relationship

Cloud vertical distribution and thickness grows with increased SST (CloudSat and CALIPSO)



Climate Change and the Monsoon – Tropical Ocean warming

Heat gain is uneven among earth system components
— Ocean gaining more than 90% of the heat



Oceans are a big deep reservoir of a liquid with high heat capacity.

Almost all (more than 90%) of Earth's heat gain goes into the oceans.

Rate is 8×10^{21} J/yr, or 2.5×10^{14} J/sec.

≈ 4 Hiroshima atomic bomb detonation per second

(1 detonation = 6.3×10^{13} Joules

So, about 2 billion atomic bombs since 1998)


Atmospheric heat gain is an especially small part of the total.

Atmosphere + Land + Ice accounts for less than 10% of the heat gain.

Climate Change and the Monsoon – Tropical Ocean warming

Heat gain in the ocean is also uneven


— Indian and Atlantic Oceans have warmed up




Climate Change and the Monsoon – Tropical Ocean warming

Surface warming uneven through different periods
— monotonous warming in the Indian and Atlantic Oceans


HADISST 1901–1950 JJAS trend per year



HADISST 1950–2012 JJAS trend per year




HADISST 1979–2012 JJAS trend per year



Climate Change and the Monsoon – Tropical Ocean warming

Subsurface warming in the Indian Ocean

— Increased heat transport from the Pacific to Indian Ocean




Indian Ocean heat content has increased abruptly, which accounts for more than 70% of the global ocean heat gain in the upper 700 m during the past decade.

Climate Change and the Monsoon – Tropical Ocean warming


— western Indian Ocean warmed up to 1.2°C

SST trend




Basin-wide warming, with significant warming over western Indian Ocean.

SST Climatology




Monotonous warming over west nullifies zonal SST gradient

Time-series [1901–2012]



Changes in SST seasonal cycle




SST change largest in summer


Climate Change and the Monsoon – Tropical Ocean warming

WIO: Pacific-Indian Ocean teleconnection during summer — Walker Circulation


(a) Correlation: east Pacific SSTa vs Global SSTa, June-Sept mean



Neutral / La Nina conditions




El Nino conditions



Climate Change and the Monsoon – Tropical Ocean warming

Asymmetry in ENSO forcing: Influence of El Niño > La Niña




Climate Change and the Monsoon – Tropical Ocean warming

Skewness in ENSO forcing:

Increase in Frequency and Magnitude of El Niños


SST anomalies [°C]: east Pacific vs WIO, June-Sept mean




Detrended anomalies show increase in frequency and strength of El Niños. The warm events over Indian Ocean also has increased. Occasionally, they cross the El Niño criteria (1 S.D. = 0.77 degC).

Positive skewness over east Pacific in recent decades >>

(a) SST Skewness [1901-1950]



(b) SST Skewness [1951-2012]




Climate Change and the Monsoon – Tropical Ocean warming

Ideally,


Increased ocean warming = more rainfall

Increased land-sea contrast = more rainfall

Increased ocean warming




Increased land-sea thermal contrast




Climate Change and the Monsoon – Tropical Ocean warming

but it's a weak South Asian Monsoon




Decreasing trend in precipitation from Pakistan through central India to Bangladesh.
Significant over central Indian subcontinent (horse-shoe pattern)




Climate Change and the Monsoon – Tropical Ocean warming

EOF₄ corresponds to ocean warming trend

EOFs and corresponding PCs of precipitation




Correlation of SST with PC




Climate Change and the Monsoon – Tropical Ocean warming

Indian Ocean warming well correlated with weak Precip.

(a) Trend in IMD Precip.




(b) Trend in CRU Precip.




(a) & (b)

Decreasing trend in precipitation from Pakistan through central India to Bangladesh. Significant over central Indian subcontinent (horse-shoe pattern)

(c) Correlation: WIO HadISST vs IMD Precip.



(d) Correlation: WIO ERSST vs CRU Precip.




(c) & (d)

Trend and correlation with western Indian Ocean warming has similar patterns!


Climate Change and the Monsoon – Tropical Ocean warming

Land-sea thermal contrast over South Asian domain


Indian Ocean-large warming, Subcontinent-suppressed warming



Increased SST results in intense vertical development of convection




(c) Trend in land-sea temperature difference



Climate Change and the Monsoon – Tropical Ocean warming


Weakening local Hadley circulation:
Convection enhanced over ocean and suppressed over land

Observations: trend in vertical velocity (1948–2012)



WIO warming extends the warm pool,
and increases ocean convection

Large scale upward motion over the Indian ocean
(10S–10N), extending up to the upper troposphere
and favoring intense local convection.




Compensated by subsidence of air over the
subcontinent (10–20°N), inhibiting convection
over the landmass and drying the region.


Climate Change and the Monsoon – Tropical Ocean warming

Weakened Monsoon precip/winds due to warming Model simulations with Indian Ocean warming

Model simulated warming of WIO




Model simulated response to warming



Competition between ocean and land rainfall:
SST warming extends the warm pool, increases ocean rainfall
...but results in decreased rainfall over the subcontinent
– horseshoe pattern in model simulations with increased IO warming


Model simulated vertical velocity in response to Indian Ocean warming




Climate Change and the Monsoon – Tropical Ocean warming

Other Mechanisms

Rise in Indo-Pacific SST increases tropical western Pacific monsoon rainfall, which incites a Rossby wave that forces descending air to the west, drying South Asia.



Climate Change and the Monsoon – Tropical Ocean warming




Future?

CMIP5 future projections suggest further warming of the Indian Ocean. Will the monsoon decrease further?

These future projections also suggest increasing monsoon rainfall (Sharmila et al 2015).


However it is to be noted that these models fail to reproduce the present day monsoon (Sabeerali et al 2014, Saha et al 2014)

Climate Change and the Monsoon – Tropical Ocean warming




CMIP5 models fail to reproduce the Indian Ocean SSTs
–bias in thermocline/equatorial dynamics?

Observations Vs. CMIP5 SST in the Indian Ocean



Climate Change and the Monsoon – Tropical Ocean warming

Does the Atlantic warming has a role?



Climate Change and the Monsoon – Tropical Ocean warming


Indian Ocean warming may dampen the El Niño

Table 1 Summary of the numerical experiments with their main characteristics, including length, nudging domain and SST climatology used for the nudging in the Indian or Atlantic oceans decoupled experiments


Name	REF	FTIC	FTIC-obs	FTAC	FTAC-obs
Correction area	None	Indian Ocean 30°E–120°E 25°S–30°N	Indian Ocean 30°E–120°E 25°S–30°N	Atlantic Ocean 100°W–20°E 25°S–25°N	Atlantic Ocean 100°W–20°E 25°S–25°N
Smoothing area	None	30°S–25°S	30°S–25°S	30°S–25°S 25°N–30°N	30°S–25°S 25°N–30°N
SST data	None	REF	AVHRR	REF	AVHRR
Time duration (year)	210	110	50	110	50

Suppressing the Indian Ocean SST variability increased the ENSO strength.

Suppressed Indian Ocean variability




Suppressed Atlantic Ocean variability



Climate Change and the Monsoon – Tropical Ocean warming

Indian Ocean warming may shorten the El Niño cycle




Suppressing the Indian Ocean SST variability increased the El Niño decaying period.
i.e. increased Indian Ocean SST variability kills an El Niño at an earlier state.

Involve modulations of the surface winds in the western equatorial Pacific, which trigger eastward-propagating oceanic Kelvin waves responsible for the turnabout of ENSO (through changes in the thermocline).


Climate Change and the Monsoon – Tropical Ocean warming

Increased variance in ISO and changes in ocean-atmosphere interaction

Trend in MISO variance, 1979–2010




Changes in SST–precipitation relationship



Atmosphere takes more time to respond to a SST anomaly in the recent decade, and therefore, ISO gets more time to build up its amplitude. The increased moisture-holding capacity of the atmosphere in the recent decade due to warming and associated increase in the residence time of the atmospheric moisture may be one possible reason to increase the maximum SST lead in the recent decade.


Slowdown in the northward propagation of ISO

Northward propagation of rainfall anomalies,
1979–1988 and 2001–2010




Northward propagation of ISO is proportional to the easterly vertical wind shear and moisture gradient (Jiang et al. 2004)

Decreasing wind shear and moisture gradient over the monsoon domain might be slowing down the ISO.



Climate Change and the Monsoon – Tropical Ocean warming

Changes in Monsoon ISO



Multidecadal variability of the Monsoon

— Atlantic Multidecadal Oscillation (AMO)

Positive AMO phase → Strong NAO

→ Changes in Jetstream and storm tracks

→ Positive TT anomalies over Asia

→ Increased meridional TT gradient → Strong monsoon





Figure 7: Multidecadal oscillation of AIR (red line, obtained from 11-yr running mean of JJAS mean all India rainfall) and Atlantic multidecadal oscillation (AMO, black line). AMO is based on 60-month running mean of monthly anomalies averaged over Atlantic north of Equator.

Modeling the Monsoon – weather prediction

Weather forecast skills in the northern and southern hemispheres



The convergence of the curves for Northern Hemisphere (NH) and Southern Hemisphere (SH) after 1999 indicates the breakthrough in exploiting satellite data.

Modeling the Monsoon

Two kinds of atmospheric predictability

Initial conditions (IC) refer to constraints imposed on the solution in **time**.

In other words, the solutions of a time-dependent differential equation must have a specific value at a certain point in time.

(current state of atmospheric quantities based on satellite and in-situ observations)

Boundary conditions (BC) refer to constraints imposed on the solution in **space**.

In other words, the solutions of a time-dependent differential equation must have a specific value at a certain point in space.

(greenhouse gases – CO₂, aerosols, albedo, SSTs, Sea Ice, Land Surface – vegetation etc.)

Predictability of the first kind focuses on the initial value problem:


how uncertainties in the initial state of the climate system amplify and spoil the forecast at a given lead time. This is the dominant source of error in weather forecasting. Errors in the atmospheric analysis grow such that the current predictability horizon is often of the order of a week or so.

Predictability of the second kind focuses on the boundary value problem:

how predictable changes in the boundary conditions that affect climate can provide predictive power. A common class of second-kind predictability studies use atmosphere models with prescribed sea surface temperatures (SSTs) in order to assess the upper limit of predictability associated with a perfect knowledge of the future ocean state.


Modeling the Monsoon

Two kinds of atmospheric predictability



Modeling the Monsoon – weather prediction


Initial value problem – forecast uncertainty



Probabilities indicate the percent of ensemble members that predict the mean precipitation

Modeling the Monsoon – weather prediction


Physical processes of importance to weather prediction



These processes are not explicitly resolved in current NWP models but they are represented via parameterizations describing their contributions to the resolved scales in terms of mass, momentum and heat transfers.

Modeling the Monsoon – weather prediction

Key challenge areas





Advances in forecast skill will come from scientific and technological innovation in computing, the representation of physical processes in parameterizations, coupling of Earth-system components, the use of observations with advanced data assimilation algorithms, and the consistent description of uncertainties through ensemble methods and how they interact across scales.

Modeling the Monsoon – weather prediction

Convection Schemes ...


Convection schemes calculates the tendencies of temperature and humidity by the result of interaction between grid scale circulation and sub-grid scale cloud ensemble. Cloud liquid water and cumulus cloud fraction is also determined for radiative processes.



Modeling the Monsoon – weather prediction


Convection Schemes ...

http://climate.snu.ac.kr/gcmdocu/Phy_Cum.htm – Check this website for code!




Modeling the Monsoon – weather prediction

Super-parameterized Convection



Modeling the Monsoon – weather prediction




Super-parameterized Convection



Modeling the Monsoon – weather prediction


Super-parameterized Climate Forecast System (SP-CFS)

Convective tendencies are explicitly simulated with a **Cloud Resolving Model** running in each GCM grid column which replaces the traditional cumulus parameterization of the GCM.






Modeling the Monsoon – weather prediction

Super-parameterized Climate Forecast System (SP-CFS)



Improvement in tropospheric temperature bias is seen in TT gradient. Even though the Gradient looks reasonable in both CFS and SPCFS, but the bias is seen when we see the North and South boxes individually. The TT-gradient in CFS perhaps is consistent with reasonable circulation pattern but deficient moisture, leading to dry monsoon.

Right result due to wrong reason in CFSv2?




Short Range Ensemble Prediction System at IITM

GEFS T574 and GFS T1534 use Revised simplified Arakawa Schubert convective parameterization scheme. (Han and Pan 2011, Ganai et al. 2015)

– improvement of diurnal cycle of convection. The major difference between the new and old schemes lies in the heating and cooling behavior in lower-atmospheric layers above the planetary boundary layer


SCHEMATIC OF GEFS (SL) T574 L64 RUNNING AT IITM



Modeling the Monsoon – weather prediction

Extended Range Ensemble Prediction System at IITM

Ensemble perturbations: At ECMWF, initial perturbations are generated using singular–vector technique (by determining error growth) and an unperturbed control run (Buizza and Palmer, 1995). At NCEP, the ensemble of initial perturbations are generated in a similar way as at ECMWF, but breeding vectors (Toth and Kalnay, 1993) are used instead of singular vectors. At Meteorological Service of Canada (MSC) different combinations of parameterization schemes is used to generate ensemble of initial conditions for their medium range prediction (Houtekamer et al., 1996, 2005)




Modeling the Monsoon – weather prediction



Extended Range Ensemble Prediction System at IITM

Bias correction technique

SST Bias from Long Simulation




SST Bias
20 day Lead




Lead
Dependant
Bias

Abhilash et al., 2013




Northward propagation of ISO



Extended Range Ensemble Prediction System at IITM


The ensembles are generated not only by perturbing the initial condition, but also by using different resolutions, parameters, and coupling configurations of the same model (CFS and its atmosphere component, the Global Forecast System). Each of these configurations was created to address the role of different physical mechanisms known to influence error growth on the 10–20-day time scale. MME has been formulated using 21 ensembles of GFSbc, 11 Ensembles of CFS126 and 11 ensembles of CFS382.

Prediction of 2015 monsoon:




Modeling the Monsoon – weather prediction

Prediction of heavy rainfall events



Uttarakhand event in June 2013


Evolution of Potential Vorticity (PV; $\times 10^{-7} \text{ s}^{-1}$) anomalies at 700 hPa and mean sea level pressure



Modeling the Monsoon – seasonal prediction

Improvement with enhanced resolution and improved coupling

GFDL Coupled Model East Pacific Precipitation (150°W-90°W)




Observed rainfall

GFDL CM2.1
2° Atmosphere
1° Ocean

GFDL CM2.3
1° Atmosphere
1° Ocean

GFDL CM2.4
1° Atmosphere
1/4° Ocean

GFDL CM2.5
1/2° Atmosphere
1/4° Ocean



Modeling the Monsoon – seasonal prediction

Improvement with enhanced resolution and improved coupling





Figure: Gabriel Vecchi


Modeling the Monsoon – seasonal prediction

Dynamical model prediction skills

Dynamical AGCM prediction skill,
CC = 0.39




Dynamical CGCM prediction skill,
NCEP CFSv2: CC=0.45



Modeling the Monsoon – seasonal prediction

Monsoon Mission – CFSv2



CFSv2 model biases and improvement National Monsoon Mission

Major Biases in CFSv2

Dry bias over India

Cold bias in SST

Mixed layer bias

Cold land and trop. Temperature

Excess Eurasian snow

Excessive convective rainfall over tropics

(Roxy et al. 2012, Saha et al. 2013)

Attempts to reduce these biases:

Convective parameterization (New SAS, Han and Pan, 2011; Ganai et al. 2014)

Cloud Microphysics (Hazra et al. 2015; Abhik et al. 2016)

Super parameterization (Goswami et al. 2015)

Improved snow physics in Land Surface Model (Saha et al. 2016)


High resolution model (Ramu et al. 2015)

Earth System Model (Swapna et al. 2014)

Modeling the Monsoon – seasonal prediction

CFSv2 model biases and improvement

5-AGCM EM hindcast skill (21Yr)




AGCMs were unable to predict ASM rainfall.

Inherent lack of ocean-atmospheric coupling, and treating monsoon as a slave to prescribed SST results in the failure.

Earlier version models

1979-1999




Poor skill

High skill

Preeti et al., (2009)

Latest models (ENSEMBLES)

1979-1999



Improved skill



High skill

Rajeevan et al., (2011)

Modeling the Monsoon – seasonal prediction


CFSv2 model biases and improvement

T382L64 Skill of Rainfall/SST




Modeling the Monsoon – seasonal prediction

CFSv2 model biases and improvement



Monsoon Mission Model Performance (Prediction Skill as well as interannual variance) is better than other models for Indian Monsoon.




Modeling the Monsoon – seasonal prediction


predictable mode analysis

ocean–atmospheric processes – 47% of total variance

EOFs and corresponding PCs of precipitation




Correlation of SST with PC



Modeling the Monsoon – seasonal prediction

— models unable to account for changing ocean-atmospheric processes and teleconnections due to recent global warming


Despite enormous progress in predicting ISMR since 1886, the operational forecasts during recent decades (1989–2012) have little skill. This recent failure is largely due to the models' inability to capture new predictability sources emerging during recent global warming, such as ENSO Modoki.



Modeling the Monsoon – seasonal prediction

— models unable to reproduce ENSO cycle and magnitude

Correlations Nino34 SST (12-1) SST - Year +1



Modeling the Monsoon – climate prediction

Seasonal forecast model to a climate model

THE IITM EARTH SYSTEM MODEL

Transformation of a Seasonal
Prediction Model to a Long-Term
Climate Model

BY P. SWAPNA, M. K. ROXY, K. APARNA, K. KULKARNI, A. G. PRAJEESH,
K. ASHOK, R. KRISHNAN, S. MOORTHY, A. KUMAR, AND B. N. GOSWAMI

Better ocean physics and representation of mean state of ocean – SST, MLD and dynamics

Improved Energy Balance in IITM ESMv2: Reduction of positive net radiation bias

Bias of OLR is reduced largely by including TKE dissipative heating (Bretherton et al. 2012)


Replace of Zhou –Carr Cloud Microphysics by Brad–Ferrier Scheme

Reduction of bias in the Sea–Ice distribution


Modeling the Monsoon – climate prediction

Seasonal forecast model to a climate model

Global mean surface temperature



Annual mean SST bias





The drift in surface temperature is minimum in IITM ESMv1 compared to CFSv2

Significant reduction in cold SST bias in tropical IO and subtropical Pacific

Modeling the Monsoon – climate prediction




Seasonal forecast model to a climate model



ENSO is better.

Modeling the Monsoon – climate prediction

Seasonal forecast model to a climate model



ENSO – Monsoon relation is better.